

---

# ОРГАНИКАЛЫҚ ХИМИЯ

# ОРГАНИЧЕСКАЯ ХИМИЯ

# ORGANIC CHEMISTRY

UDC 547.7/.8:547.796.1

O.A. Nurkenov<sup>1</sup>, S.D. Fazylov<sup>1</sup>, T.M. Seilkhanov<sup>2</sup>, A.E. Arinova<sup>1,3</sup>,  
A.Zh. Issayeva<sup>1,3</sup>, Zh.B. Satpaeva<sup>1,3</sup>, G.Zh. Karipova<sup>1</sup>

<sup>1</sup>*Institute of Organic Synthesis and Coal Chemistry of the Republic of Kazakhstan, Karaganda, Kazakhstan;*

<sup>2</sup>*Sh. Ualikhanov Kokshetau State University, Kazakhstan;*

<sup>3</sup>*Ye.A. Buketov Karaganda State University, Kazakhstan  
(E-mail: nurkenov\_oral@mail.ru)*

## Synthesis and structure of condensed biheterocycles on the basis of 3-amino-1,2,4-triazole

The article is devoted to the development of preparatively convenient methods for the synthesis of new derivatives of 3-amino-1,2,4-triazole with the aim of constructing new condensed systems of industrially important substances. The synthesis of triazolopyrimidines, prepared by three-component condensation of 1,3-dicarbonyl compounds (anilide of acetoacetic acid and acetoacetic ether) with substituted aromatic aldehydes and 3-amino-1,2,4-triazole. It is shown that by melting at 120–140 °C, in the absence of a solvent in equimolecular amounts of acetylacetil anilide and acetoacetic ether with a mixture of 3-amino-1,2,4-triazole and substituted aromatic aldehydes (3-ethoxy-4-hydroxybenzaldehyde, salicylic aldehyde), leads to the 7-aryl-5-methyl-N-phenyl-4,7-dihydro-[1,2,4-triazolo][1,5-a]pyrimidine-6-carboxamides and ethyl 7-aryl-5-methyl-4,7-dihydro-[1,2,4-triazolo][1,5-a]pyrimidin-6-carboxylate-titrates respectively. The structures of the synthesized compounds were studied by <sup>1</sup>H and <sup>13</sup>C NMR spectroscopy, as well as with the data of the two-dimensional spectra of COSY (<sup>1</sup>H-<sup>1</sup>H) and HMQC (<sup>1</sup>H-<sup>13</sup>C). The values of chemical shifts, multiplicity and integrated intensity of the <sup>1</sup>H and <sup>13</sup>C signals in one-dimensional NMR spectra are determined. Homo- and heteronuclear interactions were established using spectra in the formats COSY (<sup>1</sup>H-<sup>1</sup>H) and HMQC (<sup>1</sup>H-<sup>13</sup>C), confirming the structure of the compounds under study.

**Keywords:** 3-amino-1,2,4-triazole, 7-aryl-5-methyl-N-phenyl-4,7-dihydro-[1,2,4-triazolo][1,5-a]pyrimidine-6-carboxamides, aromatic aldehydes, <sup>1</sup>H- and <sup>13</sup>C NMR-spectra.

### *Introduction*

The growing needs of mankind in new materials predetermine the vector orientation and tasks of organic synthesis, which involve the development of fundamentally new, convenient, economically and technologically sound, safe and competitive synthesis methods, as well as the production of previously unknown chemical products. One of the modern trends in organic synthesis is the using of the polycomponent condensations. Their attractiveness for synthetic chemists is connected with the combinatorial possibilities, the environmentally oriented idea of «green chemistry», one-pot and the rejection of the solvent as the preferred methods of practical implementation, allowing the synthesis of extensive libraries of substances in a short time. Multicomponent reactions involving 1,3-dicarbonyl compounds (1,3-DCC), aldehydes, mono- and bi-N-nucleophiles play an important role in fine organic synthesis and medical chemistry as universal methods for the synthesis of a large number of biologically active compounds, for example, derivatives of pyridines, pyrimidines [1, 2].

Pyrimidines are an important component of nucleic acids, and they have been used as building blocks in pharmaceutical preparations for the synthesis of antiviral [3], anticancer [4], antibacterial and antifungal agents [5].

Previously [6], we synthesized the condensed biheterocycles — tetrazolopyrimidines obtained by three-component condensation of anilide of acetoacetic acid with substituted aromatic aldehydes and the monohydrate of 5-aminotetrazole. The closest structural analogue of 5-aminotetrazole is 3-amino-1,2,4-triazole. It should be noted that derivatives of 1,2,4-triazole have wide range of physiological activity: hepatoprotective, wound healing, antiviral, germicidal, etc. [7, 8]. They are used as additives to high photographic materials, exhibit fungicidal, herbicidal and insecticidal properties [9, 10].

Continuing research into the synthesis of pyrimidine compounds [11, 12] with the aim of constructing new condensed systems, we carried out the reaction of substituted aromatic aldehydes with 3-amino-1,2,4-triazole and 1,3-dicarbonyl compounds (anilide of acetoacetic acid and acetoacetic acid ether), leading to the formation of triazolopyrimidines.

### Experimental

The  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of compounds (1, 2) were recorded in DMSO-d6 using a JNN-ECA 400 spectrometer (400 and 100 MHz on  $^1\text{H}$  and  $^{13}\text{C}$  nuclei) from Jeol, Japan. The survey was carried out at room temperature using a DMSO solvent. Chemical shifts are measured relative to signals of residual protons or carbon atoms of a deuterated solvent.

*General procedure for the preparation of triazolopyrimidines (1–4).* A mixture of 1,3-dicarbonyl compound, aromatic aldehyde and 3-amino-1,2,4-triazole in equimolar amounts was held at 130–150 °C for 5–10 minutes until gas evolution ceased. The reaction mixture was cooled to room temperature, treated with ethyl alcohol, and the precipitate was filtered off. The precipitate was recrystallized from acetonitrile.

*Ethyl 7-(3-Ethoxy-4-hydroxyphenyl)-5-methyl-N-phenyl-4,7-dihydro-[1,2,4]triazolo[1,5-a]pyrimidine-6-carboxamide (1).* The yield of the product (1) was 2.16 g (56 %), m.p. 245–247 °C.

*7-(2-Hydroxyphenyl)-5-methyl-N-phenyl-4,7-dihydro-[1,2,4]triazolo[1,5-a]pyrimidine-6-carboxamide (2).* Product yield (2) was 2.6 g (63 %) with m.p. 100–105 °C.


*Ethyl 7-(3-ethoxy-4-hydroxyphenyl)-5-methyl-N-phenyl-4,7-dihydro-[1,2,4]triazolo[1,5-a]pyrimidine-6-carboxylate (3).* The yield of product (3) was 3.55 g (43 %), m.p. 225–228 °C.


*Ethyl 7-(2-hydroxyphenyl)-5-methyl-4,7-dihydro-[1,2,4]triazolo[1,5-a]pyrimidine-6-carboxylate (4).* The yield of the product (4) was 3 g (83 %), m.p. 145–148 °C.

### Results and discussion

In this paper we present the results of the research of the product of a three-component reaction of the anilide of acetoacetic acid and acetoacetic ether with substituted aromatic aldehydes and 3-amino-1,2,4-triazole in the absence of a solvent and a catalyst. The three-component Biginelli reaction, originally proposed for the synthesis of dihydropyrimidine-2(1*H*)-one by the reaction of aromatic aldehyde, urea and acetoacetic ether, is now used with great success in the synthesis of dihydropyrimidine-2(1*H*)-thione and various annelated pyrimidines [13–23].

We found that the fusion of equimolecular amounts of an acetoacetyl anilide with a mixture of 3-amino-1,2,4-triazole and substituted aromatic aldehydes at 130–150 °C resulted in 7-aryl-5-methyl-N-phenyl-4,7-dihydro[1,2,4-triazolo][1,5-a]pyrimidine-6-carboxamides (1, 2), and with acetoacetic ether — ethyl 7-aryl-5-methyl-4,7-dihydro-[1,2,4-triazolo[1,5-a]pyrimidine-6-carboxylate (3, 4). The resulting compounds (1–4) are pale yellow powders soluble in DMF, DMSO, heated in ethanol, acetonitrile, insoluble in water.






Based on the analysis of the published data [4–6, 11, 12], it is possible to assume that at the first stage of the reaction, as a result of the interaction of the 1,3-dicarbonyl compound with the aromatic aldehyde molecule, an unsaturated compound I is formed, the interaction of which with 3-amino-1,2,4-triazole, leads to the final reaction product. It is shown that in the 3-amino-1,2,4-triazole molecule there are two unequal nitrogen atoms in positions 2 and 4 of the heterocycle, which allows us to consider the two most probabilistic reaction products. However, the main and nucleophilic character of nitrogen in position 2 of the heterocycle, apparently, causes the reaction to proceed in the direction predominantly to form the structure of A.

The proposed mechanism is confirmed by analysis of the IR and NMR spectra of <sup>1</sup>H-, <sup>13</sup>C. The IR spectra of compounds (1, 2) exhibit bands due to stretching vibrations of amide groups (1672–1684 cm<sup>-1</sup>) and NH groups (3085–3120 cm<sup>-1</sup>).

The <sup>1</sup>H NMR spectrum of compound (1) is characterized by the presence in the strong-field region of the spectrum at 1.19 ppm triplet signal with an intensity of 3H with <sup>3</sup>J 6.9 Hz atoms of the methyl group CH<sub>3</sub><sup>21</sup> ethyl radical. Another CH<sub>3</sub><sup>18</sup> methyl radical bound to the C<sup>8</sup> atom of a heterocyclic six-membered ring



does not have spin-spin interactions with neighboring protons and is manifested at 2.12 ppm singlet signal with a content of 3H. In this part of the spectrum at 2.46 and 3.32 ppm the presence of solvent signals of DMSO-d6 and residual amounts of water was noted. The methylene protons of the hydroxyethyl radical were resonated with a multiplet signal in the range 3.76–3.89 ppm with an integrated intensity of 2H. The singlet signal with intensity IH at 6.39 ppm corresponds to proton H<sup>6</sup>. The protons of the aromatic system H<sup>15</sup> and H<sup>14</sup>, which have hydrogen atoms adjacent to each other, which contribute to the splitting of the spectrum, were manifested by doublet signals at 6.56 (<sup>3</sup>J 7.8 Hz) and 6.56 (<sup>3</sup>J 8.2 Hz) ppm with intensities IH. The proton H<sup>11</sup> of the aromatic cycle under consideration,

which does not have hydrogen atoms adjacent to it, was indicated by a singlet at 6.71 ppm with an intensity of 1H. For an unsubstituted phenyl radical, all the protons appeared in the aromatic zone of the spectrum. The asymmetric proton H<sup>27</sup> was manifested in the form of a triplet signal at 6.96 ppm, corresponding to the intensity IH, with <sup>3</sup>J 7.3 Hz. Equivalent H<sup>26,28</sup> protons were manifested by triplet signals at 7.20 ppm with <sup>3</sup>J 7.8 Hz with an integrated intensity of 2H, whereas equivalent protons of H<sup>25,29</sup> were detected by doublet signals at 7.48 ppm with <sup>3</sup>J 7.9 Hz with a 2H content. The multiplicity of the signals under consideration, their integrated intensity, and the values of the spin-spin interaction constants are in good agreement with the NMR indices for aromatic systems. The methyl proton H<sup>5</sup> of the five-membered heterocyclic system adjacent to two nitrogen atoms was manifested by the expected singlet at 7.59 ppm with an integrated intensity of 1H. The protons of the hydroxyl group H<sup>22</sup> and the protons bound to the nitrogen atoms of H<sup>23</sup> and H<sup>9</sup> were manifested in the form of singlets with an integrated intensity of 1H each at 8.90, 9.66 and 10.10 ppm respectively.

In the  $^{13}\text{C}$  NMR spectrum of compound (1), the carbon atoms of the methyl groups resonated at 15.15 ( $\text{C}^{21}$ ) and 17.81 ( $\text{C}^{18}$ ) ppm in the strong-field part of the spectrum. Then they showed themselves at 60.47 ppm  $\text{C}^6$  atoms and at 64.43 ppm  $\text{C}^{20}$  atoms, which have a small screening effect on the carbon nucleus. The other carbons of the bicyclic fragment were resonated at 104.35 ( $\text{C}^7$ ), 136.75 ( $\text{C}^8$ ), 148.24 ( $\text{C}^3$ ), and 150.23 ( $\text{C}^5$ ) ppm. Signals with chemical shifts at 113.35 ( $\text{C}^{11}$ ), 115.96 ( $\text{C}^{14}$ ), 120.02 ( $\text{C}^{25,29}$ ), 120.25 ( $\text{C}^{15}$ ), 123.77 ( $\text{C}^{27}$ ), 129.08 ( $\text{C}^{26,28}$ ), 132.17 ( $\text{C}^{10}$ ), 139.58 ( $\text{C}^{24}$ ), 146.86 ( $\text{C}^{12}$ ) and 147.29 ( $\text{C}^{13}$ ) ppm belong to the carbon nuclei of the two aromatic rings. In the most weakly-field region of the spectrum at 165.58 ppm the carbonyl carbon atom  $\text{C}^{16}$  resonated.

The structure of compound (1) was also confirmed by two-dimensional spectroscopy of COSY ( $^1\text{H}$ - $^1\text{H}$ ) NMR spectroscopy and HMQC ( $^1\text{H}$ - $^{13}\text{C}$ ), which makes it possible to establish spin-spin interactions of homo- and heteronuclear nature (Fig. 1–4). The observed correlations in the molecule are shown in the diagram. In the spectra of the  $^1\text{H}$ - $^1\text{H}$  COSY compound, spin-spin correlations are observed through three proton bonds of neighboring methylene groups  $\text{H}^{25,29}$ - $\text{H}^{26,28}$  (7.47, 7.20 and 7.17, 7.48),  $\text{H}^{26}$ - $\text{H}^{27}$  (7.20, 6.96 and 6.96, 7.21) and  $\text{H}^{27}$ - $\text{H}^{28}$  (6.96, 7.20 and 7.20, 6.96). The coordinates 3.84, 1.18 and 1.16, 3.80 correspond to homolytic interaction through three bonds of neighboring aliphatic protons  $\text{H}^{20}$  and  $\text{H}^{21}$ . Heteronuclear interactions of protons with carbon atoms through one bond were established using  $^1\text{H}$ - $^{13}\text{C}$  HMQC spectroscopy for all pairs present in the compound:  $\text{H}^5$ - $\text{C}^5$  (7.57, 150.23),  $\text{H}^{25,29}$ - $\text{C}^{25,29}$  (7.48, 119.93),  $\text{H}^{26,28}$ - $\text{C}^{26,28}$  (7.19, 129.15),  $\text{H}^{27}$ - $\text{C}^{27}$  (6.94, 123.73),  $\text{H}^{11}$ - $\text{C}^{11}$  (6.70, 113.34),  $\text{H}^{20}$ - $\text{C}^{20}$  (3.82, 64.45),  $\text{H}^{14}$ - $\text{C}^{14}$  (6.65, 120.32),  $\text{H}^6$ - $\text{C}^6$  (6.37, 60.47),  $\text{H}^{18}$ - $\text{C}^{18}$  (2.11, 17.83) and  $\text{H}^{21}$ - $\text{C}^{21}$  (1.17, 15.05).




Figure 1. Correlations of COSY ( $^1\text{H}$ - $^1\text{H}$ ) compound (1)






Figure 2. Correlations of HMQC ( $^1\text{H}$ - $^{13}\text{C}$ ) compound (1)

Figure 3. COSY ( $^1\text{H}$ - $^1\text{H}$ ) NMR spectrum of compound (1)Figure 4. HMQC ( $^1\text{H}$ - $^{13}\text{C}$ ) NMR spectrum of compound (1)

The structures of other compounds (2), (3), and (4) are also confirmed by single-dimensional  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectroscopy (Table) and homo- and heteronuclear correlations of COSY and HMQC NMR.

T a b l e

 $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopy data for compounds (1–4)

Compound No.	$\delta$ , ppm.	
	$^1\text{H}$	$^{13}\text{C}$
1	1.19 t (3H, $\text{CH}_3^{14}$ , $^3\text{J}$ 6.9 Hz), 2.12 s (3H, $\text{CH}_3^{18}$ ), 3.76–3.89 m (2H, $\text{CH}_2^{20}$ ), 6.39 s (1H, $\text{H}^6$ ), 6.56 d (1H, $\text{H}^{15}$ , $^3\text{J}$ 7.8 Hz), 6.65 d (1H, $\text{H}^{14}$ , $^3\text{J}$ 8.2 Hz), 6.71 s (1H, $\text{H}^{11}$ ), 6.96 t (1H, $\text{H}^{27}$ , $^3\text{J}$ 7.3 Hz), 7.20 t (2H, $\text{H}^{26,28}$ , $^3\text{J}$ 7.8 Hz), 7.48 d (2H, $\text{H}^{25,29}$ , $^3\text{J}$ 7.8 Hz), 8.90 s (1H, $\text{OH}^{22}$ ), 9.66 s (1H, $\text{NH}^{23}$ ). 10.10 s (1H, $\text{NH}^9$ )	15.15 ( $\text{C}^{21}$ ), 17.81 ( $\text{C}^{18}$ ), 60.47 ( $\text{C}^6$ ), 64.43 ( $\text{C}^{20}$ ), 104.35 ( $\text{C}^7$ ), 113.35 ( $\text{C}^{11}$ ), 115.96 ( $\text{C}^{14}$ ), 120.02 ( $\text{C}^{25,29}$ ), 120.25 ( $\text{C}^{15}$ ), 123.77 ( $\text{C}^{27}$ ), 129.08 ( $\text{C}^{26,28}$ ), 132.17 ( $\text{C}^{10}$ ), 136.75 ( $\text{C}^8$ ), 139.58 ( $\text{C}^{24}$ ), 146.86 ( $\text{C}^{12}$ ), 147.29 ( $\text{C}^{13}$ ), 148.24 ( $\text{C}^3$ ), 150.23 ( $\text{C}^5$ ), 165.58 ( $\text{C}^{16}$ )
2	2.26 s (3H, $\text{CH}_3^{18}$ ), 5.78 s (1H, $\text{H}^3$ ), 6.70 t (1H, $\text{H}^{13}$ ), 6.95 d (1H, $\text{H}^{12}$ ), 7.15–7.26 m (3H, $\text{H}^{14,24,26}$ ), 7.36–7.59 m (3H, $\text{H}^{22,23,25}$ ), 7.87 d (1H, $\text{H}^{15}$ ), 9.64 s (1H, $\text{OH}^{19}$ ), 9.99 s (1H, $\text{NH}^6$ ). 10.37 s (1H, $\text{NH}^{20}$ )	19.20 ( $\text{C}^{18}$ ), 60.29 ( $\text{C}^3$ ), 118.86 ( $\text{C}^{22,26}$ ), 119.18 ( $\text{C}^{12}$ ), 119.71 ( $\text{C}^{13}$ ), 120.19 ( $\text{C}^{24}$ ), 128.06 ( $\text{C}^{10}$ ), 128.81 ( $\text{C}^{15}$ ), 129.02 ( $\text{C}^{14}$ ), 129.09 ( $\text{C}^{23,25}$ ), 129.32 ( $\text{C}^{21}$ ), 150.18 ( $\text{C}^5$ ), 151.22 ( $\text{C}^9$ ), 152.38 ( $\text{C}^7$ ), 156.43 ( $\text{C}^{21}$ ), 170.88 ( $\text{C}^{16}$ )
3	1.01 t (3H, $\text{CH}_3^{25}$ , $^3\text{J}$ 7.0 Hz), 1.24 t (3H, $\text{CH}_3^{21}$ , $^3\text{J}$ 7.0 Hz), 2.35 s (3H, $\text{CH}_3^{18}$ ), 3.87–3.98 m (4H, $\text{CH}_2^{20}$ , $\text{CH}_2^{22}$ ), 6.11 s (1H, $\text{H}^6$ ), 6.50 dd (1H, $\text{H}^{15}$ , $^3\text{J}$ 8.2, 1.8 Hz), 6.64 d (1H, $\text{H}^{14}$ , $^3\text{J}$ 8.2 Hz), 6.72 s (1H, $\text{H}^{11}$ ), 6.96 t (1H, $\text{H}^{27}$ , $^3\text{J}$ 7.3 Hz), 7.20 t (2H, $\text{H}^{26,28}$ , $^3\text{J}$ 7.8 Hz), 7.48 d (2H, $\text{H}^{25,29}$ , $^3\text{J}$ 7.8 Hz), 8.90 s (1H, $\text{OH}^{22}$ ), 9.66 s (1H, $\text{NH}^{23}$ ). 10.10 s (1H, $\text{NH}^9$ )	14.49 ( $\text{C}^{25}$ ), 15.21 ( $\text{C}^{21}$ ), 18.88 ( $\text{C}^{18}$ ), 59.62 ( $\text{C}^6$ ), 59.82 ( $\text{C}^{24}$ ), 64.41 ( $\text{C}^{20}$ ), 98.08 ( $\text{C}^7$ ), 113.41 ( $\text{C}^{11}$ ), 115.91 ( $\text{C}^{14}$ ), 119.80 ( $\text{C}^{15}$ ), 133.70 ( $\text{C}^{10}$ ), 146.70 ( $\text{C}^{3,8}$ ), 147.09 ( $\text{C}^{12}$ ), 147.41 ( $\text{C}^{13}$ ), 150.45 ( $\text{C}^5$ ), 165.75 ( $\text{C}^{16}$ )
4	1.16 t (3H, $\text{CH}_3^{22}$ , $^3\text{J}$ 6.9 Hz), 2.26 s (3H, $\text{CH}_3^{18}$ ), 3.99 k (2H, $\text{CH}_2^{21}$ , $^3\text{J}$ 7.0 Hz), 6.76 d (1H, $\text{H}^{12}$ , $^3\text{J}$ 8.8 Hz), 6.80 s (1H, $\text{H}^3$ ), 6.93 t (1H, $\text{H}^{13}$ , $^3\text{J}$ 6.3 Hz), 7.06 t (1H, $\text{H}^{14}$ , $^3\text{J}$ 7.3 Hz), 7.68 d (1H, $\text{H}^{15}$ , $^3\text{J}$ 8.2 Hz), 9.38 s (1H, $\text{OH}^{19}$ ), 9.65 s (1H, $\text{NH}^6$ )	14.32 ( $\text{C}^{22}$ ), 19.18 ( $\text{C}^{18}$ ), 44.87 ( $\text{C}^3$ ), 61.06 ( $\text{C}^{21}$ ), 98.67 ( $\text{C}^4$ ), 120.01 ( $\text{C}^{12}$ ), 122.79 ( $\text{C}^{13}$ ), 125.44 ( $\text{C}^{10}$ ), 127.07 ( $\text{C}^{15}$ ), 128.87 ( $\text{C}^{14}$ ), 147.60 ( $\text{C}^9$ ), 150.20 ( $\text{C}^5$ ), 151.19 ( $\text{C}^7$ ), 156.42 ( $\text{C}^{11}$ ), 165.39 ( $\text{C}^{16}$ )

## Conclusions

The studies carried out for the first time the synthesis of novel 7-aryl-5-methyl-N-phenyl-4,7-dihydro-[1,2,4-triazolo][1,5-a]pyrimidine-6-carboxyethylamides and 7-aryl-5-methyl-4,7-dihydro-[1,2,4-triazolo][1,5-a]pyrimidine-6-carboxylates. On basis of analysis of spectral data of their structure and methods NMR  $^1\text{H}$ - and  $^{13}\text{C}$ -spectroscopy as well as two-dimensional spectra data COSY ( $^1\text{H}$ - $^1\text{H}$ ) and HMQC ( $^1\text{H}$ - $^{13}\text{C}$ ) provided a mechanism of the reaction whereby the formation of the final product takes place through the intermediate step of reacting 1,3-dicarbonyl compounds with molecules of aromatic aldehyde.

*The source of research funding.* The work was carried out with the financial support of the Committee of Science and the Ministry of Education of the Republic of Kazakhstan on «Program-Targeted Financing», No. Registration 0115PK01782.

## References

- Крыльский Д.В. 5-Амино-4-фенилпиразол в реакциях трехкомпонентной конденсации / Д.В. Крыльский, Х.С. Шихалиев, А.С. Чувашлев // Журн. орг. химии. — 2010. — Т. 46, Вып. 3. — С. 416–422.
- Семенов В.Э. Взаимодействие производных 6-метилурацила с ацетилацетоном и ацетоуксусным эфиром / В.Э. Семенов, Е.С. Крылова, И.В. Галыметдинова, Д.Р. Шарафутдинова, В.Д. Акамсин, В.С. Резник // Журн. общ. химии. — 2010. — Т. 80, Вып. 7. — С. 1192–1197.
- Гетероциклические соединения / под ред. Р. Эльдерфилда. — М.: Мир, 1969. — Т. 8. — С. 8.
- Гейн В.Л. Синтез 7-арил-N-метил(N, N-диэтил)-5-метил-4,7-дигидротетразоло-[1,5-а]пиримидин-6-карбоксамидов / В.Л. Гейн, Т.М. Замараева, Н.В. Носова, М.И. Вахрин, П.А. Слепухин // Журн. орг. химии. — 2012. — Т. 48, Вып. 3. — С. 422–425.

- 5 Гейн В.Л. Синтез и взаимодействие с гидразингидратом метиловых эфиров 7-арил-6-(2-тиеноил)-4,7-дигидротетразоло-[1,5-*a*]пиримидин-5-карбоновых кислот / В.Л. Гейн, В.В. Мишунин, Е.П. Цыплякова, М.И. Вахрин, П.А. Слепухин // Журн. орг. химии. — 2011. — Т. 47, Вып. 7. — С. 1060–1064.
- 6 Нуркенов О.А. Синтез 7-арил-5-метил-N-фенил-4,7-дигидротетразоло[1,5-*a*]пиримидин-6-карбоксамидов / О.А. Нуркенов, С.Д. Фазылов, Т.М. Сейлханов, А.Е. Аринова, Ж.Б. Сатпаева, М.З. Мулдахметов, А.Ж. Исаева, Г.Ж. Карипова, А.Б. Мукашев // Изв. НАН РК. Сер. химии и технологии. — 2017. — № 1(421). — С. 76–81.
- 7 Пат. 2370492. Российская Федерация. Способ получения морфолиния 3-метил-1,2,4-триазолил-5-тиоацетата, проявляющего гепатозащитную, ранозаживляющую и противовирусную активность / Е.Г. Кныш, Н.И. Трубчанин. Опубл. 20.01.00.
- 8 Овсепян Т.Р. Синтез замещенных 1,2,4-триазолов и 1,3,4-тиадиазолов / Т.Р. Овсепян, Э.Р. Диланян, А.П. Евгоян // Химия гетероциклических соединений. — 2004. — № 9. — С. 1377–1381.
- 9 Veerasha-Sharmaa P.M. Novel Synthesis of 5-Substituted-3-phenylindole-2-(1,2,4-triazole) Derivatives / P.M. Veerasha-Sharmaa, M.G. Purohit // Synth. Commun. — 2008. — Vol. 38, No. 9. — P. 1381–1388.
- 10 Zheng-Jun Quan. Copper-catalyzed click synthesis of functionalized 1,2,3-triazoles with 3,4-dihydropyrimidinone or amide group via a one-pot four-component reaction Zheng-Jun Quan, Qiong Xu, Zhang Zhang, Yu-Xia Da, Xi-Cun Wang // Tetrahedron Lett. — 2013. — Vol. 69, No. 38. — P. 881–887.
- 11 Фазылов С.Д. Трехкомпонентная циклоконденсация тиомочевины, ацетоуксусного эфира и замещенных бензальдегидов в условиях микроволновой активации / С.Д. Фазылов, А.Е. Аринова, О.А. Нуркенов, А.В. Болдашевский // Журн. общ. химии. — 2012. — Т. 82, Вып. 2. — С. 343–343.
- 12 Нуркенов О.А. Синтез и модификация 5-этокси-6-метил-4-(4-диэтиламинофенил)-2-тиоксо-1,2,3,4-тетрагидропирамидин-5-карбоксилата / О.А. Нуркенов, С.Д. Фазылов, Т.М. Сейлханов, Т.С. Животова, А.Е. Аринова, Ж.Б. Сатпаева, А.Ж. Исаева, Г.Ж. Карипова, А.Б. Мукашев, З.М. Мулдахметов // Изв. НАН РК. Сер. химии и технологии. — 2016. — № 2(416). — С. 77–84.
- 13 Вдовина С.В. Новые возможности классической реакции Биджинелли / С.В. Вдовина, А.О. Мамедов // Успехи химии. — 2008. — Т. 77, Вып. 2. — С. 1091–1128.
- 14 Suresh T. Synthesis and bronchodilatory activity of new 4-aryl-3,5-bis(2-chlorophenyl)-carbamoyl-2,6-dimethyl-1,4-dihydropyridines & their 1-substituted analogues / T. Suresh, S.K. Swamy, V.M. Reddy // Ind. Journal of Chem. Sect. B. — 2007. — Vol. 46B, No. 1. — P. 115–121.
- 15 Desai B.G. Synthesis and QSAR studies of 4-substituted phenyl-2,6-dimethyl-3,5-bis-N-(substituted phenyl)carbamoyl-1,4-dihydropyridines as potential antitubercular agents / B.G. Desai, D. Sureja, Y. Nalapara, A. Shah, A.K. Saxena // Bioorg. Med. Chem. — 2001. — № 9. — P. 1993.
- 16 Shah A. Advanced Dihydropyridines as Novel Multidrug Resistance Modifiers and Reversing Agents / A. Shah, J. Bariwal, J. Molnár, M. Kawase, N. Motohashi // Topics in Heterocyclic Chemistry. — 2008. — Vol. 15. — P. 201–252.
- 17 Десенко С.М. Азагетероциклы на основе ароматических непредельных кетонов / С.М. Десенко, В.Д. Орлов. — Харьков: Фолио, 1998. — 148 с.
- 18 Wipf P. A solid phase protocol of the Biginelli dihydropyrimidine synthesis suitable for combinatorial chemistry / P. Wipf, V. Cunningham // Tetrahedron Lett. — 1995. — № 36. — P. 7819.
- 19 Gupta R. Synthesis of 3,4-dihydropyrimidin-2(1H)-ones / R. Gupta, A.K. Gupta, S. Paul, P.L. Kachroo // Ind. Journal of Chem. Sect. B. — 1995. — № 34. — P. 151–153.
- 20 Grover G.J. Pharmacologic Profile of the Dihydropyrimidine Calcium Channel Blockers SQ 32,547 and SQ 32,946 / G.J. Grover, S. Dzwonczyk, D.M. McMullin, C.S. Normandin, S.J. Moreland // Journal Cardiovasc. Pharmacol. — 1995. — № 26. — P. 289–291.
- 21 Колосов М.А. 3-N-Ацилирование 5-карбетокси-6-метил-4-фенил-3,4-дигидро-пиримидин-2-она в системе карбоновая кислота-SOCl<sub>2</sub> / М.А. Колосов, В.Д. Орлов // Химия гетероциклических соединений. — 2005. — № 2. — С. 292–293.
- 22 Kappe C.O. 100 Years of the Biginelli dihydropyrimidine synthesis / C.O. Kappe // Tetrahedron Lett. — 1993. — Vol. 49, No. 32. — P. 6937–6963.
- 23 Колосов М.А. Синтез производных 3-метил-4-фенил-3а,4,5,6,7а-гексагидро-1Н-пиразоло[4,5-*d*]пиримидин-6-она / М.А. Колосов, В.Д. Орлов // Химия гетероциклических соединений. — 2007. — № 10. — С. 1586–1588.

О.А. Нуркенов, С.Д. Фазылов, Т.М. Сейлханов, А.Е. Аринова,  
А.Ж. Исаева, Ж.Б. Сатпаева, Г.Ж. Карипова

### 3-Амино-1,2,4-триазол негізінде конденсирленген бигетероциклдердің синтезі мен құрылымы

Макала 3-амино-1,2,4-триазолдың жана туындыларын синтездеуде препаратты тиімді әдістерін дамытып, өндіріске қажетті жаңа конденсирленген жүйелерді құру мақсатына арналған. 1,3-ди-карбонилді қосылыстардың (ацетосірке қышқылының анилиді және ацетосірке қышқылы), орынауыстырылған ароматтық альдегидтер және 3-амино-1,2,4-триазолымен үшкомпонентті конденсациясы арқылы алынатын триазоло-пиримидиндерді синтездеу туралы мәліметтер көлтірілді. Еріткішсіз эквимолекулярлық мөлшерде ацетосірке қышқылының анилиді мен ацетосірке қышқылын қоспасымен 3-амино-1,2,4-триазол және орынауыстырылған ароматтық альдегидтер (3-этокси-4-гидроксибензальдегид, салицил альдегид) қоспасымен 120–140 °C-да балқыту арқылы сәйкесінше

7-арил-5-метил-N-фенил-4,7-дигидро-[1,2,4-триазоло][1,5-*a*]пиридин-6-карбоксамидтер мен этил 7-арил-5-метил-4,7-дигидро-[1,2,4-триазоло][1,5-*a*]пиридин-6-карбоксилаттар альнуы көрсетілді. Синтезделген қосылыстардың құрылымы ЯМР  $^1\text{H}$ - мен  $^{13}\text{C}$ -спектроскопия, екі кеңістікті COSY ( $^1\text{H}$ - $^1\text{H}$ ) және HMQC ( $^1\text{H}$ - $^{13}\text{C}$ ) әдістерімен зерттелді. Бір кеңістікті ЯМР спектрлерінде химиялық қозғалыстардың сандық мәндері, мультиплеттілігі және  $^1\text{H}$ ,  $^{13}\text{C}$  сигналдарының интегралды карқындылығы анықталды. Зерттелген қосылыстардың құрылымы COSY ( $^1\text{H}$ - $^1\text{H}$ ) и HMQC ( $^1\text{H}$ - $^{13}\text{C}$ ) спектрлері көмегімен гомо- и гетероядролық аракатынасын анықтау арқылы дәлелденді.

*Кілт сөздер:* 3-амино-1,2,4-триазол, 7-арил-5-метил-N-фенил-4,7-дигидро-[1,2,4-триазоло][1,5-*a*]пиридин-6-карбоксамидтер, ароматтық альдегидтер, ЯМР  $^1\text{H}$ - и  $^{13}\text{C}$ -спектрлер.

О.А. Нуркенов, С.Д. Фазылов, Т.М. Сейлханов, А.Е. Аринова,  
А.Ж. Исаева, Ж.Б. Сатпаева, Г.Ж. Карипова

## Синтез и строение конденсированных бигетероциклов на основе 3-амино-1,2,4-триазола

Статья посвящена разработке препаративно удобных способов синтеза новых производных 3-амино-1,2,4-триазола с целью построения новых конденсированных систем промышленно важных веществ. Приведены данные по синтезу триазолопиридинов, полученных трехкомпонентной конденсацией 1,3-дикарбонильных соединений (анилид ацетоуксусной кислоты и ацетоуксусный эфир) с замещенными ароматическими альдегидами и 3-амино-1,2,4-триазолом. Показано, что сплавление при 120–140 °C в отсутствие растворителя эквимолекулярных количеств анилида ацетоуксусной кислоты и ацетоуксусного эфира со смесью 3-амино-1,2,4-триазола и замещенных ароматических альдегидов (3-этокси-4-гидроксибензальдегид, салициловый альдегид) приводит к 7-арил-5-метил-N-фенил-4,7-дигидро-[1,2,4-триазоло][1,5-*a*]пиридин-6-карбоксамидам и этил 7-арил-5-метил-4,7-дигидро-[1,2,4-триазоло][1,5-*a*]пиридин-6-карбоксилатам соответственно. Исследовано строение синтезированных соединений методами ЯМР  $^1\text{H}$ - и  $^{13}\text{C}$ -спектроскопии, а также данными двумерных спектров COSY ( $^1\text{H}$ - $^1\text{H}$ ) и HMQC ( $^1\text{H}$ - $^{13}\text{C}$ ). Определены значения химических сдвигов, мультиплетность и интегральная интенсивность сигналов  $^1\text{H}$  и  $^{13}\text{C}$  в одномерных спектрах ЯМР. С помощью спектров в форматах COSY ( $^1\text{H}$ - $^1\text{H}$ ) и HMQC ( $^1\text{H}$ - $^{13}\text{C}$ ) установлены гомо- и гетероядерные взаимодействия, подтверждающие структуру исследуемых соединений.

**Ключевые слова:** 3-амино-1,2,4-триазол, 7-арил-5-метил-N-фенил-4,7-дигидро-[1,2,4-триазоло][1,5-*a*]пиридин-6-карбоксамиды, ароматические альдегиды, ЯМР  $^1\text{H}$ - и  $^{13}\text{C}$ -спектры.

## References

- 1 Krylskii, D.V., Shikhaliev, Kh.S., & Chuvashlev, A.S. (2010). 5-Amino-4-fenilpirazol v reaktsiakh trekhkomponentnoy kondensatsii [Three-component condensations with 5-amino-4-phenylpyrazole]. *Zhurnal orhanicheskoy khimii — Journal of Organic Chemistry*, 46, 416–422 [in Russian].
- 2 Semenov, V.E., Krylova, E.S., Galyametdinova, I.V., Sharafutdinova, D.R., Akamsin, V.D., & Reznik, V.S. (2010). Vzaimodeystviye proizvodnykh 6-metiluratsila s atsetilatsetonom i atsetouksusnym efironom [Reaction of 6-methyluracyl derivatives with acetylacetone and ethyl acetoacetate]. *Zhurnal obshchey khimii — Russian Journal of General Chemistry*, 80, 1192–1197 [in Russian].
- 3 Elderfield R. (Eds.). (1969) *Heterotsiklicheskie soyedineniya* [Heterocyclic compounds]. Moscow: Mir [in Russian].
- 4 Gein, V.L., Zamaraeva, T.M., Nosov, N.V., Vakhruin, M.I., & Slepukhin, P.A. (2012). Sintez 7-aryl-N-metil (N,N-diethyl)-5-metil-4,7-dihidrotetrazolo-[1,5-*a*]pirimidin-6-karboksamidov [Synthesis of 7-aryl-N-methyl(N,N-diethyl)-5-methyl-4,7-dihydro-tetrazolo[1,5-*a*]pyrimidine-6-carboxamides]. *Zhurnal orhanicheskoy khimii — Journal of Organic Chemistry*, 48, 422–425 [in Russian].
- 5 Gein, V.L., Mishunin, V.V., Tsyplyakovka, E.P., Vakhruin, M.I., & Slepukhin, P.A. (2011). Sintez i vzaimodeystviye s hidrazinhydratom metilovykh efirov 7-aryl-6-(2-tiienoil)-4,7-dihidrotetrazolo-[1,5-*a*]pirimidin-5-karbonovykh kislot [Synthesis and reaction with 7-aryl-6-(2-thienoyl)-4,7-dihidrotetrazolo[1,5-*a*]pyrimidine-5-carboxylic acid methyl ester methyl ester]. *Zhurnal orhanicheskoy khimii — Journal of Organic Chemistry*, 47, 1060–1064 [in Russian].
- 6 Nurkenov, O.A., Fazylov, S.D., Seilkhanov, T.M., Arinova, A.E., Satpaeva, Zh.B., & Muldakhmetov, M.Z., et al. (2017). Sintez 7-арил-5-метил-N-фенил-4,7-дигидро-тетразоло[1,5-*a*]пиридин-6-карбоксамидов [Synthesis of 7-aryl-5-methyl-N-phenyl-4,7-dihydrotetrazolo[1,5-*a*]pyrimidine-6-carboxamides]. *Izvestiya Natsionalnoi Akademii Nauk Respubliki Kazakhstan. Seriya khimii i tekhnologii — Proceedings of National Academy of Sciences of Kazakhstan. Series of Chemistry and Technology*, 76–81 [in Russian].
- 7 Knyshev E.G., Trubchanin N.I. (Publ. 20.01.2000). Patent of Russian Federation, No. 2370492. Sposob polucheniya morfoliniya 3-metil-1,2,4-triazolil-5-tioatsetata, proyavlyayushcheho hepatozashchitnyu, ranozazhivlyayushchuyu i protivovirusnuyu aktivnost' [Method for the preparation of morpholinium of 3-methyl-1,2,4-triazolyl-5-thioacetate exhibiting hepatoprotective, wound healing and antiviral activity]. [in Russian].

- 8 Hovsepyan, T.R., Dilanyan, E.R., & Evgonyan, A.P. (2004). Sintez zameshchennykh 1,2,4-triazolov i 1,3,4-tiadiazolov [Synthesis of substituted 1,2,4-triazoles and 1,3,4-thiadiazoles]. *Khimiya heterotsiklichesikh soyedineniy — Chemistry of Heterocyclic Compounds*, 40, 1377–1381 [in Russian].
- 9 Veerasha-Sharmaa, P.M., & Purohitb, M.G. (2008). Novel Synthesis of 5-Substituted-3-phenylindole-2-(1,2,4-triazole) Derivatives. *Synth. Commun.*, 38, 1381–1388.
- 10 Zheng-Jun, Quan, Qiong, Xu, Zhang, Zhang, Yu-Xia, Da, & Xi-Cun, Wang. (2013). Copper-catalyzed click synthesis of functionalized 1,2,3-triazoles with 3,4-dihydropyrimidinone or amide group via a one-pot four-component reaction. *Tetrahedron*, 69, 881–887.
- 11 Fazylov, S.D., Arinova, A.E., Nurkenov, O.A., & Boldashevsky, A.B. (2012). Trekhkomponentnaya tsiklokondensatsiya tiromocheviny, atsetouksusnoho efira i zameshchennykh benzal'dehidov v usloviyah mikrovolnovoy aktivatsii [Three-component cyclocondensation of thiourea, acetoacetic ether and substituted benzaldehydes under microwave activation conditions]. *Zhurnal obshchey khimii — Russian Journal of General Chemistry*, 82, 343–343 [in Russian].
- 12 Nurkenov, O.A., Fazylov, S.D., Seilkhanov, T.M., Zhivotova, T.S., Arinova, A.E., & Issayeva, A.Zh. et al. (2016). Sintez i modifikatsiya 5-etoksi-6-metil-4-(4-diethylaminofenil)-2-tiokso-1,2,3,4-tetrahidropirimidin-5-karboksilata [Synthesis and modification of 5-ethoxy-6-methyl-4-(4-diethylaminophenyl)-2-thioxo-1,2,3,4-tetrahydropyrimidine-5-carboxylate]. *Izvestiya Natsional'noy Akademii Nauk Respubliki Kazakhstan. Seriya khimii i tekhnologii — Proceedings of National Academy of Sciences of Kazakhstan. Series of Chemistry and Technology*, 77–84 [in Russian].
- 13 Vdovina, S.V., & Mamedov, A.O. (2008). Novyye vozmozhnosti klassicheskoy reaktsii Bidzhinelli [New possibilities of classical Biginelli reaction]. *Uspekhi khimii — Russian Chemical Reviews*, 1091–1128 [in Russian].
- 14 Suresh, T., Swamy, S.K., & Reddy, V.M. (2007). Synthesis and bronchodilatory activity of new 4-aryl-3,5-bis(2-chlorophenyl)-carbamoyl-2,6-dimethyl-1,4-dihydropyridines & their 1-substituted analogues. *Ind. Journal of Chem. Sect. B.*, 46B, 1, 115–121.
- 15 Desai, B.G., Sureja, D., Nalapara, Y., Shah, A., & Saxena, A.K. (2001). Synthesis and QSAR studies of 4-substituted phenyl-2,6-dimethyl-3,5-bis(N-substituted phenyl)carbamoyl-1,4-dihydropyridines as potential antitubercular agents. *Bioorg. Med. Chem.*, 1993.
- 16 Shah, A., Bariwal, J., Molnár, J., Kawase, M., & Motohashi, N. (2008). Advanced Dihydropyridines as Novel Multidrug Resistance Modifiers and Reversing Agents. *Topics in Heterocyclic Chemistry*, 15, 201–252.
- 17 Desenko, S.M., & Orlov, V.D. (1998). *Azaheterotsikly na osnove aromaticheskikh nepredel'nykh ketonov [Azaheterocycles based unsaturated aromatic ketones]*. Kharkov: Folio [in Russian].
- 18 Wipf, P., & Cunningham, V. (1995). A solid phase protocol of the Biginelli dihydropyrimidine synthesis suitable for combinatorial chemistry. *Tetrahedron Lett.*, 7819.
- 19 Gupta, R., Gupta, A.K., Paul, S., & Kachroo, P.L. (1995). Synthesis of 3,4-dihydropyrimidin-2(1H)-ones. *Ind. Journal of Chem. Sect. B.*, 151–153.
- 20 Grover, G.J., Dzwonczyk, S., McMulltn, D.M., Normandin, C.S., & Moreland, S.J. (1995). Pharmacologic Profile of the Dihydropyrimidine Calcium Channel Blockers SQ 32,547 and SQ 32,946. *Journal Cardiovasc. Pharmacol.*, 289–291.
- 21 Kolosov, M.A., & Orlov, V.D. (2005). 3-N-Atsilirovaniye 5-karbetski-6-metil-4-fenil-3,4-dihidro-pirimidin-2-ona v sisteme karbonovoy kislotoy-SOCl<sub>2</sub> [3-N-Acylation of 5-Ethoxycarbonyl-6-methyl-4-phenyl-3,4-dihydropyrimidin-2-one in the carboxylic acid-SOCl<sub>2</sub> system]. *Khimiya heterotsiklichesikh soyedineniy — Chemistry of Heterocyclic Compounds*, 41, 292–293 [in Russian].
- 22 Kappe, C.O. (1993). 100 Years of the Biginelli dihydropyrimidine synthesis. *Tetrahedron*, 49, 6937–6963.
- 23 Kolosov, M.A., & Orlov, V.D. (2007). Sintez proizvodnykh 3-metil-4-fenil-3a,4,5,6,7a-heksahidro-1N-pirazolo[4,5-d]pirimidin-6-ona [Synthesis of derivatives of 3-methyl-4-phenyl-3a,4,5,6,7a-hexahydro-1H-pyrazolo-[4,5-d]pyrimidin-6-one]. *Khimiya heterotsiklichesikh soyedineniy — Chemistry of Heterocyclic Compounds*, 47, 1586–1588 [in Russian].